

Eighteen (18) Statistically Significant studies showing an association of Abortion with subsequent Very PreTerm Birth, and Abortion with Very Low Birth Weight

- 1 Reime B, Schuecking BA, Wenzlaff P. Reproductive Outcomes in Adolescents Who Had a Previous Birth or an Induced Abortion Compared to Adolescents' First Pregnancies. *BMC Pregnancy and Childbirth* 2008;8:4
- 2+ Voigt M, Olbertz D, Fusch C, Krafczyk D, Briese V, Schneider KT. The influence of previous pregnancy terminations, miscarriages, and still-birth on the incidence of babies with low birth weight and premature births as well as somatic classification of newborns. *Z Geburtshilfe Neonatol* 2008;212:5-12
- 3 Smith GCS, Shah I, White IR, Pell JP, Crossley JA, Dobbie R. Maternal and biochemical predictors of spontaneous preterm birth among nulliparous women: a systematic analysis in relation to degree of prematurity. *International J Epidemiology* 2006;35(5): 1169-1177
- 4 Stang P, Hammond AO, Bauman P. Induced Abortion Increases the Risk of Very Preterm Delivery; Results from a Large Perinatal Database. *Fertility Sterility*. Sept 2005;S159 [Study only published as an abstract]
- 5+ Moreau C, Kaminski M, Ancel PY, Bouyer J, et al. Previous induced abortions and the risk of very preterm delivery: results of the EPIPAGE study. *British J Obstetrics Gynaecology* 2005;112(4): 430-437 [abstract online: www.blackwell-synergy.com/links/doi/10.1111/j.1471-0528.2004.00478.x/abs/]
- 6 Ancel PY, Lelong N, Papiernik E, Saurel-Cubizolles MJ, Kaminski M. History of induced abortion as a risk factor for preterm birth in European countries: results of EUROPOP survey. *Human Reprod* 2004;19(3):734-740.
- 7+ Ancel PY, Saurel-Cubizolles M-J, Renzo GCD, Papiernik E, Breart G. Very and moderate preterm births: are the risk factors different? *British J Obstetrics Gynaecology* 1999;106:1162-1170.
- 8+ Zhou W, Sorenson HT, Olsen J. Induced Abortion and Subsequent Pregnancy Duration. *Obstetrics & Gynecology* 1999;94:948-953.
- 9+ Martius JA, Steck T, Oehler MK, Wulf K-H. Risk factors associated with preterm (<37+0 weeks) and early preterm (<32+0 weeks): univariate and multi-variate analysis of 106 345 singleton births from 1994 statewide perinatal survey of Bavaria. *European J Obstetrics*

Gynecology Reproductive Biology 1998;80:183-189.

- 10+ Lumley J. The association between prior spontaneous abortion, prior induced abortion and preterm birth in first singleton births. *Prenat Neonat Med* 1998;3:21-24.
- 11+ Lumley J. The epidemiology of preterm birth. *Bailliere's Clin Obstet Gynecology* 1993;7(3):477-498
- 12+ Algert C, Roberts C, Adelson P, Frammer M. Low birth weight in New South Wales, 1987: a Population-Based Study. *Aust New Zealand J Obstet Gynaecol* 1993;33:243-248
- 13+* Zhang J, Savitz DA. Preterm Birth Subtypes among Blacks and Whites. *Epidemiology* 1992;3:428-433.
- 14+ Mueller-Heubach E, Guzik DS. Evaluation of risk scoring in a preterm birth prevention study of indigent patients. *Amer J Obstetrics & Gynecol* 1989;160:829-837.
- 15+ Lumley J. Very low birth-weight (less than 1500g) and previous induced abortion: Victoria 1982-1983. *Aust NZ J Obstet Gynecol* 1986;26:268-272.
- 16 Schuler D, Klinger A. Causes of low birth weight in Hungary. *Acta Paediatrica Hungarica* 1984;24:173-185
- 17+ Levin A, Schoenbaum S, Monson R, Stubblefield P, Ryan K. Association of Abortion With Subsequent Pregnancy Loss. *JAMA* 1980;243(24):2495-2499
- 18 Van Der Slikke JW, Treffers PE. Influence of induced abortion on gestational duration in subsequent pregnancies. *BMJ* 1978; 1:270-272 [$>95\%$ confident of preterm risk for gestation less than 32.0 weeks].

-
- * studies that included spontaneous and induced abortions (SIA) but did not report PTB/LBW risk separately for each
+ studies that found dose/response (the more SIAs, the higher the risk)
-